

Andrew Parker, MD 1120 Raintree Circle, Suite 280 Allen, TX 75013 Phone: (214) 383 - 9356 Fax: (214) 383 - 9886

# REHABILITATION FOLLOWING ARTHROSCOPIC ANTERIOR SHOULDER LABRAL REPAIR

Wear shoulder brace with abduction pillow 24/7 except in shower, when working with PT, or doing ROM exercises at home for first 6 weeks postoperatively.

Phase I – Immediate Postoperative Phase "Restrictive Motion" (Weeks 0-6)

Goals: Protect the anatomic repair

Prevent negative effects of immobilization Promote dynamic stability and proprioception

Diminish pain and inflammation

#### Weeks 0-2

- Elbow/hand ROM
- Hand gripping exercises
- Passive and gentle active assistive ROM exercise
  - Flexion to 70 degrees week 1
  - Flexion to 90 degrees week 2
  - ER/IR with arm 30 degrees abduction
    - ER to 5-10 degrees
    - IR to 45 degrees
    - \*\*NO active ER or Extension or Abduction
- Submaximal isometrics for shoulder musculature
- Rhythmic stabilization drills ER/IR
- Proprioception drills
- Cryotherapy, modalities as indicated

## Weeks 3-4

- Continue gentle ROM exercises (PROM and AAROM)
  - Flexion to 125 degrees at end of week 3, 145 at week 4
  - Abduction to 90 degrees
  - ER/IR at 45 degrees abd in scapular plane
  - ER in scapular plane to 20-25 degrees
  - IR in scapular plane to 45-55 degrees
  - \*\*NOTE: Rate of progression based on evaluation of the patient
- No excessive ER, extension or elevation
- Continue isometrics and rhythmic stabilization (submax)
- · Core stabilization program
- Initiate scapular strengthening program
- Continue use of cryotherapy

#### Weeks 5-6

Gradually improve ROM



Andrew Parker, MD 1120 Raintree Circle, Suite 280 Allen, TX 75013 Phone: (214) 383 - 9356 Fax: (214) 383 - 9886

- Flexion to 145 degrees
- ER at 45 degrees abduction: 55-50 degrees
- IR at 45 degrees abduction: 55-60 degrees
- May initiate stretching exercises
- Initiate exercise tubing ER/IR (arm at side)
- Scapular strengthening
- PNF manual resistance

#### II. Phase II – Intermediate Phase: Moderate Protection Phase (Weeks 7-14)

Goals: Gradually restore full ROM (week 10)

Preserve the integrity of the surgical repair

Restore muscular strength and balance
Enhance neuromuscular control

#### Weeks 7-9

- Gradually progress ROM;
  - Flexion to 170 degrees
  - Initiate ER/IR at 90 degrees abd
  - ER at 90 degrees abduction: 70-80 degrees at week 7
  - ER to 90 degrees at weeks 8-9
  - IR at 90 degrees abduction: 60-70 degrees
- Continue to progress isotonic strengthening program
- Continue PNF strengthening

# Weeks 10-14

- · May initiate slightly more aggressive strengthening
- Progress isotonic strengthening exercises
- Continue all stretching exercises
  - \*\*Progress ROM to functional demands (i.e. overhead athlete)
- Progress to isotonic strengthening (light and restricted ROM)

## III. Phase III – Minimal Protection Phase (Week 15-20)

Goals: Maintain full ROM

Improve muscular strength, power and endurance

Gradually initiate functional activities

#### Criteria to Enter Phase III

- 1. Full non-painful ROM
- 2. Satisfactory stability
- 3. Muscular strength (good grade or better)
- 4. No pain or tenderness

## Weeks 15-18

- Continue all stretching exercises (capsular stretches)
- Continue strengthening exercises:



Andrew Parker, MD 1120 Raintree Circle, Suite 280 Allen, TX 75013 Phone: (214) 383 - 9356 Fax: (214) 383 - 9886

- Throwers ten program or fundamental exercises
- PNF manual resistance
- Endurance training
- Restricted sport activities (light swimming, half golf swings)
- Initiate interval sport program week 16-18

## Weeks 18-20

- Continue all exercise listed above
- Process interval sport program (throwing, etc.)

# IV. Phase IV – Advanced Strengthening Phase (Weeks 21-24)

٧.

Goals: Enhance muscular strength, power and endurance Progress functional activities

Maintain shoulder mobility

#### **Criteria to Enter Phase IV**

- 1. Full non-painful ROM
- 2. Satisfactory static stability
- 3. Muscular strength 75-80% of contralateral side
- 4. No pain or tenderness

#### Weeks 21-24

- Continue flexibility exercises
- Continue isotonic strengthening program
- NM control drills
- Plyometric strengthening
- Progress interval sport programs

# V. Phase V – Return to Activity Phase (Months 6-9)

Goals: Gradual return to sport activities

Maintain strength, mobility and stability

#### Criteria to Enter Phase V

- 1. Full functional ROM
- 2. Satisfactory isokinetic test that fulfills criteria
- 3. Satisfactory shoulder stability
- 4. No pain or tenderness

## **Exercises**

- Gradually progress sport activities to unrestrictive participation
- Continue stretching and strengthening program